

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

B94

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification</b> <sup>6</sup> : <b>C12N 15/12, 15/85, 5/10, C07K 14/47, A61K 38/17</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 97/00319</b> <b>(43) International Publication Date:</b> 3 January 1997 (03.01.97)
<b>(21) International Application Number:</b> PCT/GB96/01388 <b>(22) International Filing Date:</b> 11 June 1996 (11.06.96) <b>(30) Priority Data:</b> 9511935.0 13 June 1995 (13.06.95) <b>GB</b> <b>(71) Applicant (for all designated States except US):</b> SMITHKLINE BEECHAM PLC [GB/GB]; New Horizons Court, Brentford, Middlesex TW8 9EP (GB). <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> BROWNE, Michael, Joseph [GB/GB]; (GB). CHAPMAN, Conrad, Gerald [GB/GB]; (GB). CLINKENBEARD, Helen, Elizabeth [GB/GB]; (GB). ROBINSON, Jeffrey, Hugh [GB/GB]; SmithKline Beecham Pharmaceuticals, Coldharbour Road, The Pinnacles, Harlow, Essex CM19 5AD (GB). <b>(74) Agent:</b> RUTTER, Keith; SmithKline Beecham, Corporate Intellectual Property, SB House, Great West Road, Brentford, Middlesex TW8 9BD (GB).	<b>(81) Designated States:</b> AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>	
<b>(54) Title:</b> CHIMERIC LEPTIN FUSED TO IMMUNOGLOBULIN DOMAIN AND USE		
<b>(57) Abstract</b>  Chimeric leptin which are proteins comprising leptin or a mutant or a variant thereof fused to a human immunoglobulin domain. One favoured immunoglobulin domain is the human immunoglobulin Fc domain. The chimeric derivatives of leptin have, despite their large molecular size, good pharmacological activity combined with prolonged clearance rates. These derivatives of leptin are therefore indicated to be particularly useful for the treatment or prophylaxis of obesity or diseases and conditions associated with obesity such as atherosclerosis, hypertension and type II diabetes.		

**CHIMERIC LEPTIN FUSED TO IMMUNOGLOBULIN DOMAIN AND USE**

The present invention relates to a novel compound being a novel chimeric protein, to a process for the preparation of such a compound, a pharmaceutical composition comprising such a compound and the use of such a compound in medicine, especially for the treatment of obesity and associated diseases.

European Patent Application, Publication number 0 464 533 discloses fusion proteins comprising various portions of the constant region of immunoglobulin molecules together with another human protein or part thereof. European Patent Application, Publication number 0 297 882 discloses fusion proteins comprising various portions of the plasminogen molecule with part of another human protein.

Zhang et al. (Nature: 372, 425 - 432; 1994) describe the positional cloning of a mouse obese gene and its human homologue. The sequence of the Open Reading Frame (ORF) of the mouse gene predicts a polypeptide of 167 amino acids and Zhang et al. predicted the presence of a signal sequence which would lead to the production of a mature protein of 146 residues. The human homologue was disclosed as having a similar size of 146 amino acids for the mature protein. Zhang et al. showed the presence of a primary translation product of approximate size of 18 kilodaltons (kD) with truncation to a 16kD product on addition of microsomal membranes, consistent with the production of a pre-protein and the removal of an N-terminal signal sequence. Zhang et al also disclose the potential use of the human obese gene product (hereinafter 'leptin') in the treatment of obesity.

For effective, practical treatment of obesity a particularly desirable property of an obesity agent is a clearance rate in humans commensurate with patient acceptable treatment regimens, especially regimens for injectable therapies. Zhang et al. do not disclose information relating to the clearance rate of the active molecule in either mouse or humans.

The precise mechanism of action of leptin is currently unknown, however it is considered that in order to provide the observed pharmacological effects, leptin must interact with one or more receptors in the brain.

We have now discovered certain chimeric derivatives of leptin which surprisingly, despite their large molecular size, have good pharmacological activity combined with prolonged clearance rates. These chimeric derivatives of leptin are therefore indicated to be particularly useful for the treatment or prophylaxis of obesity and for the treatment or prophylaxis of diseases and conditions associated with obesity, such as atherosclerosis, hypertension and, especially, Type II diabetes. In particular these compounds are considered to be useful for administration by injection.

component suitably comprises the CH2 and CH3 domains and the hinge region including cysteine residues contributing to inter-heavy chain disulphide bonding.

For example when the IgG component is derived from IgG4 it includes cysteine residues 8 and 11 of the IgG4 hinge region (Pinck J.R. and Milstein C., Nature 5. vol216pp941-942, 1967). Preferably the IgG4 component consists of amino acids corresponding to residues 1-12 of the hinge, 1-110 of CH2 and 1-107 of CH3 of IgG4 described by Ellison J., Buxbaum J. and Hood L., DNA vol1pp11-18, 1981. In one example of a suitable mutation in IgG4, residue 10 of the hinge (residue 241, Kabat numbering) is altered from serine (S) in the wild type to proline (P) and residue 5 of CH2 10 (residue 248, Kabat numbering) is altered from leucine (L) in the wild type to glutamate (E).

DNA polymers which encode mutants or variants of the human immunoglobulin may be prepared by site-directed mutagenesis of the cDNA which codes for the required protein by conventional methods such as those described by G. Winter *et al* in Nature 15 1982, 299, 756-758 or by Zoller and Smith 1982; Nucl. Acids Res., 10, 6487-6500, or deletion mutagenesis such as described by Chan and Smith in Nucl. Acids Res., 1984, 12, 2407-2419 or by G. Winter *et al* in Biochem. Soc. Trans., 1984; 12, 224-225 or polymerase chain reaction such as described by Mikaelian and Sergeant in Nucleic Acids Research, 1992, 20, 376.

20 When used herein 'compound of the invention' or 'compounds of the invention' relates to the above mentioned chimera.

In a further aspect, the invention provides a process for preparing a compound according to the invention which process comprises expressing DNA encoding said compound in a recombinant host cell and recovering the product.

25 The DNA polymer comprising a nucleotide sequence that encodes the compound also forms part of the invention.

The process of the invention may be performed by conventional recombinant techniques such as described in Maniatis *et. al.*, Molecular Cloning - A Laboratory Manual; Cold Spring Harbor, 1982 and DNA Cloning vols I, II and III (D.M. Glover ed., 30 IRL Press Ltd).

In particular, the process may comprise the steps of:

- i) preparing a replicable expression vector capable, in a host cell, of expressing a DNA polymer comprising a nucleotide sequence that encodes said compound;
- ii) transforming a host cell with said vector;
- 35 iii) culturing said transformed host cell under conditions permitting expression of said DNA polymer to produce said compound; and
- iv) recovering said compound.

molecule encoding a said leptin or variant and a second DNA molecule encoding a said immunoglobulin domain or fragment thereof.

5 The DNA molecules may be obtained by the digestion with suitable restriction enzymes of vectors carrying the required coding sequences or by use of polymerase chain reaction technology.

The precise structure of the DNA molecules and the way in which they are obtained depends upon the structure of the desired product. The design of a suitable strategy for the construction of the DNA molecule coding for the compound is a routine matter for the skilled worker in the art.

10 The expression of the DNA polymer encoding the compound in a recombinant host cell may be carried out by means of a replicable expression vector capable, in the host cell, of expressing the DNA polymer. The expression vector is novel and also forms part of the invention.

15 The replicable expression vector may be prepared in accordance with the invention, by cleaving a vector compatible with the host cell to provide a linear DNA segment having an intact replicon, and combining said linear segment with one or more DNA molecules which, together with said linear segment, encode the compound, under ligating conditions.

20 The ligation of the linear segment and more than one DNA molecule may be carried out simultaneously or sequentially as desired.

Thus, the DNA polymer may be preformed or formed during the construction of the vector, as desired.

25 The choice of vector will be determined in part by the host cell, which may be prokaryotic, such as *E. coli*, or eukaryotic, such as mouse C127, mouse myeloma, chinese hamster ovary, Cos1 or Hela cells, fungi e.g. filamentous fungi or unicellular yeast or an insect cell such as *Drosophila*. The host cell may also be a transgenic animal.

A preferred host cell is Cos1.

Suitable vectors include plasmids, bacteriophages, cosmids and recombinant viruses derived from, for example, baculoviruses, vaccinia or Semliki Forest virus.

30 The preparation of the replicable expression vector may be carried out conventionally with appropriate enzymes for restriction, polymerisation and ligation of the DNA, by procedures described in, for example, Maniatis *et al.*, cited above. Polymerisation and ligation may be performed as described above for the preparation of the DNA polymer. Digestion with restriction enzymes may be performed in an appropriate buffer at a temperature of 20°-70°C, generally in a volume of 50µl or less with 0.1-10µg DNA.

35

in metabolic rate or oxygen consumption. Multiple injections of the active compound - at most twice daily - over a period of a week for rodents or a month for primates, also cause a reduction in body weight and in the size of discrete adipose tissue depots.

5 Clearance rates are determined by conventional plasma assay using ob-antibodies, for example ELISA methodology.

As indicated above the compounds of the present invention have useful pharmaceutical properties, in particular anti obesity activity and also for the treatment of diseases associated with obesity, such as atherosclerosis, hypertension and, especially, Type II diabetes.

10 In use the compound will normally be employed in the form of a pharmaceutical composition in association with a human pharmaceutical carrier, diluent and/or excipient, although the exact form of the composition will depend on the mode of administration.

The active compound may be formulated for administration by any suitable route and is preferably in unit dosage form. Advantageously, the composition is suitable for 15 oral, rectal, topical, parenteral, intravenous or intramuscular administration or through the respiratory tract. Preparations may be designed to give slow release of the active ingredient.

The compositions of the invention may be in the form of tablets, capsules, sachets, vials, powders, granules, lozenges, suppositories, reconstitutable powders, or liquid 20 preparations such as oral or sterile parenteral solutions or suspensions. Topical formulations are also envisaged where appropriate.

The invention therefore further provides a pharmaceutical composition comprising a compound of the invention and a pharmaceutically acceptable carrier.

25 The dosage ranges for administration of the compounds of the present invention are those to produce the desired therapeutic effect. Dosage will generally vary with age, extent or severity of the medical condition and contraindications, if any. For example in the treatment of obesity the unit dosage can vary from less than 1mg to 300mg, but typically will be in the region of 1 to 20mg per dose, in one or more doses, such as one to six doses per day, such that the daily dosage is in the range 0.02-40mg/kg.

30 Dosages and compositions for the treatment of diseases associated with obesity such as atherosclerosis, hypertension and, especially, Type II diabetes are selected from an equivalent range to that used in the treatment of obesity.

Compositions suitable for injection may be in the form of solutions, suspensions or emulsions, or dry powders which are dissolved or suspended in a suitable vehicle prior 35 to use.

Fluid unit dosage forms are prepared utilising the compound and a pyrogen-free sterile vehicle. The compound, depending on the vehicle and concentration used, can be

The invention also provides the use of a compound of the invention in the manufacture of a medicament for treating obesity or diseases associated with obesity, such as atherosclerosis, hypertension and, especially, Type II diabetes.

As indicated above the invention also encompasses cosmetic treatments.

5 Accordingly, there is also provided a compound of the invention for use in the cosmetic treatment of human or non-human mammals.

There is also provided a method for the cosmetic treatment of a human or non-human mammal, which treatment comprises administering an effective, non-toxic amount of a compound of the invention to a human or non-human mammal in need thereof.

10 Cosmetic treatment suitably includes treatment for the improvement of body appearance, such as weight reduction treatment.

The invention also extends to a cosmetic composition, comprising a compound of the invention and a carrier therefor.

15 Compositions of the invention including cosmetic compositions are formulated using known methods, for example those described in standard text books of pharmaceuticals and cosmetics, such as Harry's Cosmeticology published by Leonard Hill Books, Remington's Pharmaceutical Sciences, the British and US Pharmacopoeias.

No unexpected toxicological effects are expected when compounds of the invention are administered in accordance with the present invention.

20 The following Examples illustrate the invention but do not limit it in any way.

CCTGAATTTCTGGGGGGACCATCAGTCTTCCTGTTCCCCCAAACCCAAGGACACTCTC  
 600  
 5 ATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGACGTGAGCCAGGAAGACCCC  
 660  
 GAGGTCCAGTTCAACTGGTACGTGGATGGCGTGGAGGTGCATAATGCCAAGACAAAGCCG  
 720  
 10 CGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTCCTGCACCAG  
 780  
 GACTGGCTGAACGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGGCCTCCCGTCATCG  
 840  
 15 ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAGCCACAGGTGTACACCCCTG  
 900  
 20 CCCCCATCCCAGGAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCCTGGTCAAAGGC  
 960  
 TTCTACCCCAGCGACATCGCCGTGGAGTGGGAGAGCAATGGGCAGCCGGAGAACAACACTAC  
 1020  
 25 AAGACCACGCCTCCCGTGCTGGACTCCGACGGATCCTTCTTCCTCTACAGCAGGCTAACC  
 1080  
 GTGGACAAGAGCAGGTGGCAGGAGGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCT  
 1140  
 30 CTGCACAACCACTACACACAGAAGAGCCTCTCCCTGTCTCTGGGTAAA  
 1188  
 35

Table 2. Amino acid sequence of leptin/IgG4 chimera, 396aa

40 1 MHWGTLCGFL WLWPYLFYVQ AVPIQKVQDD TKTLIKTIVT  
 RINDISHTQS  
 51 VSSKQKVTGL DFIPGLHPIL TLSKMDQTLA VYQQILTSMP SRNVIQISND  
 45 101 LENLRDLLHV LAFSKSCHLP WASGLETLDL LGGVLEASGY  
 STEVVALSRL  
 151 QGSLQDMLWQ LDLSPGCEK YGPPCPSCPA PEFLGGPSVF  
 LFPPKPKDTL  
 50 201 MISRTPEVTC VVVDVSQEDP EVQFNWYVDG VEVHNAKTKP  
 REEQFNSTYR



Example 2.**Construction of DNA coding for fusion protein ob 1-167/IgG4 hinge-CH2-CH3 PE variant**

5 The gene coding for a fusion protein comprising the human 'ob' protein and the Hinge-CH2-CH3 region of human IgG4 PE (a form of IgG4 mutated as below) is created by recombinant DNA technology, preferably by a two-step recombinant PCR method.

10 The cDNA coding for the complete human leptin, amino acids 1-167 (numbering as Y. Zhang, R. Proenca, M. Maffei, M. Barone, L. Leopold & J. Friedman. Nature 372: 425-432) is joined at the 3' end to the 5' end of the hinge-CH2-CH3 region of the cDNA coding for the human IgG4 (PE variant) protein, shown as amino acids 168-396 in the protein sequence below.

15 The human 'ob' gene has been prepared synthetically based on the amino acid sequence of Zhang et al, and assembled in the pcDNA3 vector. The encoded protein sequence is given in Table 2.

20 Human IgG4 heavy chain PE variant. In IgG4 PE, residue 10 of the hinge (residue 241, Kabat numbering) is altered from serine (S) in the wild type to proline (P) and residue 5 of CH2 (residue 248, Kabat numbering) is altered from leucine (L) in the wild type to glutamate (E). Angal S., King D.J., Bodmer M.W., Turner A., Lawson A.D.G., Roberts G., Pedley B. and Adair R., Molecular Immunology vol30pp105-108, 1993, describe an IgG4 molecule where residue 241 (Kabat numbering) is altered from serine to proline. This change increases the serum half-life of the IgG4 molecule.

25 The IgG4 PE variant was created using PCR mutagenesis on the synthetic human IgG4 heavy chain cDNA. The sequence of the IgG4 PE variant is described in Table 1. The residues of the IgG4 nucleotide sequence which were altered to make the PE variant are as follows:

referring to Table 1:

30 residue 322 has been altered to "C" in the PE variant from "T" in the wild type;

residue 333 has been altered to "G" in the PE variant from "A" in the wild type; and

35 residues 343-344 have been altered to "GA" in the PE variant from "CT" in the wild type.

The fusion protein was expressed transiently in Cos1 cells using the pCDN vector system, as described in International Patent Application Publication number WO 96/04388. The mature protein was exported from the cells into the culture medium and was detected by anti-leptin antibody. It was shown to have a size consistent with the predicted structure by Western blotting analysis under both reducing and nonreducing conditions.

CTGGAGAACCTCCGGGATCTTCTTCACGTGCTGGCCTTCTCTAAGAGCTGCCACTTGCCC  
 360  
 5 TGGGCCAGTGGCCTGGAGACCTTGGACAGCCTGGGGGTGTCTCTGAGGCTTCAGGCTAC  
 420  
 TCCACAGAGGTGGTGGCCCTGAGCAGGCTGCAGGGGTCTCTGCAGGACATGCTGTGGCAG  
 480  
 CTGGACCTCAGCCCCGGGTGCGAGTCCAAATATGGTCCCCCATGCCACCATGCCCAGCG  
 540  
 10 CCTGAATTTGAGGGGGGACCATCAGTCTTCCTGTTCCCCCAAACCCAAGGACACTCTC  
 600  
 ATGATCTCCCGGACCCCTGAGGTCACGTGCGTGGTGGTGGACGTGAGCCAGGAAGACCCC  
 660  
 15 GAGGTCCAGTTCAACTGGTACGTGGATGGCGTGGAGGTGCATAATGCCAAGACAAAGCCG  
 720  
 CGGGAGGAGCAGTTCAACAGCACGTACCGTGTGGTCAGCGTCCTACCGTCCTGCACCAG  
 780  
 GACTGGCTGAACGGCAAGGAGTACAAGTGAAGGTCTCCAACAAAGGCCTCCCGTCATCG  
 840  
 20 ATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAGCCACAGGTGTACACCCTG  
 900  
 CCCCCATCCCAGGAGGAGATGACCAAGAACCAGGTGAGCCTGACCTGCCTGGTCAAAGGC  
 960  
 25 TTCTACCCCAGCGACATCGCCGTGGAGTTGGAGAGCAATGGGCAGCCGGAGAACAACCTAC  
 1020  
 AAGACCACGCCTCCCGTGCTGGACTCCGACGGATCCTTCTTCCTCTACAGCAGGCTAACC  
 1080  
 GTGGACAAGAGCAGGTGGCAGGAGGGGAATGTCTTCTCATGCTCCGTGATGCATGAGGCT  
 1140  
 30 CTGCACAACCACTACACACAGAAGAGCCTCTCCCTGTCTCTGGGTAAA  
 1188

Table 4: Amino acid sequence of ob 1-167/IgG4 hinge-CH2-CH3 PE variant chimera 396aa

35

SEQ ID No: 2

1 MHWGTLCGFL WLWPYLFYVQ AVPIQKVQDD TKTLIKTIVT RINDISHTQS  
 40 51 VSSKQKVTGL DFIPGLHPIL TLSKMDQTLA VYQQILTSMP SRNVIQISND  
 101 LENLRDLLHV LAFSKSCHLP WASGLETLDL LGGVLEASGY STEVVALSRL  
 151 QGSLQDMLWQ LDLSPGCESK YGPPCPPCPA PEFEGGPSVF LFPPKPKDTL  
 45 201 MISRTPEVTC VVVDVSQEDP EVQFNWYVDG VEVHNAKTKP REEQFNSTYR  
 251 VVSVLTVLHQ DWLNGKEYKC KVSNGKLPSS IEKTISKAKG QPREPQVYTL  
 50 301 PPSQEEMTKN QVSLTCLVKG FYPSDIAVEW ESNGQPENNY KTTTPVLDS

The mature protein was exported from the cells into the culture medium and was detected by anti-leptin antibody. It was shown to have a size consistent with the predicted structure by Western blotting analysis under both reducing and nonreducing conditions.

5

**Table 5. DNA sequence of ob/IgG1 chimera 1197bp**

```

ATGCATTGGGGAACCCTGTGCGGATTCTTGTGGCTTTGGCCCTATCTTTTCTATGTCCAA
60
10 GCTGTGCCCATCCAAAAGTCCAAGATGACACCAAACCCTCATCAAGACAATTGTCACC
120
AGGATCAATGACATTTACACACGCAGTCAGTCTCCTCCAAACAGAAAGTCACCGGTTTG
15 180
GACTTCATTCTGCTGGGCTCCACCCCATCCTGACCCTGTCCAAGATGGACCAGACACTGGCA
240
20 GTCTACCAACAGATCCTCACATCGATGCCTTCCAGAAACGTGATCCAAATATCCAACGAC
300
CTGGAGAACCTCCGGGATCTTCTTCACGTGCTGGCCTTCTCTAAGAGCTGCCACTTGCCC
360
25 TGGGCCAGTGGCCTGGAGACCTTGGACAGCCTGGGGGGTGTCTCGAGGCTTCAGGCTAC
420
TCCACAGAGGTGGTGGCCCTGAGCAGGCTGCAGGGGTCTCTGCAGGACATGCTGTGGCAG
30 480
CTGGACCTCAGCCCCGGGTGCGAGCCCAAATCGGCCGACAAACTCACACATGCCCACCG
540
35 TGCCCAGCACCTGAACTCCTGGGGGGACCGTCAGTCTTCTCTTCCCCC AAAACCCAAG
600
GACACCCTCATGATCTCCCGGACCCCTGAGGTCACATGCGTGGTGGTGGACGTGAGCCAC
660
40 GAAGACCCTGAGGTCAAGTTCAACTGGTACGTGGACGGCGTGGAGGTGCATAATGCCAAG
720
ACAAAGCCGCGGGAGGAGCAGTACAACAGCACGTACCGTGTGGTCAGCGTCCTCACCGTC
45 780
CTGCACCAGGACTGGCTGAATGGCAAGGAGTACAAGTGCAAGGTCTCCAACAAAGCCCTC
840
50 CCAGCCCCCATCGAGAAAACCATCTCCAAAGCCAAAGGGCAGCCCCGAGAACCACAGGTG
900
TACACCCTGCCCCCATCCCGGGAGGAGATGACCAAGAACCAGGTCAGCCTGACCTGCCG
960

```

The cDNA encoding the full length human leptin (nucleotides 1-501) is joined at the 3' end to the 5' end of the hinge-CH2-CH3 region of the IgG1 cDNA (nucleotides 508-1203).

The two amino acid linker between the two parts of the fusion is encoded by the nucleotide sequence GGTACC (502-507). See Table 1.

5 The encoded protein sequence of the leptin/IgG1(GT) chimera is given in Table 2. Leptin 1-1 (numbering as Y. Zhang, R. Proenca, M. Maffei, M. Barone, L. Leopold & J. Friedman. Nature 372:425-432), followed by the GT linker (168-169) and IgG1 H-CH2-CH3 170-401.

10 The gene coding for the human IgG1 contains a number of nucleotide substitutions compared to the IgG1 molecule described by Ellison J.W., Berson B.J. and Hood L.E., Nucleic Acids Research vol 10 No. 13 pp4071-4079, 1982. The IgG1 nucleotides which differ from the Ellison J.W. et al published sequence and the resulting amino acid substitutions are as follows (nucleotide numbering as in table 1).

15 nucleotide 519 is "G" in this variant compared to "T" in the Ellison et al sequence (silent mutation)

nucleotides 520-522 are "GCC" in this variant compared to "TGT" in the Ellison et al sequence (resulting in substitution of Ala for Cys in this variant, amino acid 174 in table 2)

20 nucleotide 759 is "T" in this variant compared to "G" in the Ellison et al sequence (silent mutation)

25 nucleotide 924 is "G" in this variant compared to "T" in the Ellison et al sequence (resulting in substitution of Glu for Asp in this variant, amino acid 308 in table 2)

nucleotide 928 is "A" in this variant compared to "C" in the Ellison et al sequence (resulting in substitution of Met for Val in this variant, amino acid 310 in table 2)

30 nucleotide 1077 is "T" in this variant compared to "C" in the Ellison et al sequence (silent mutation)

nucleotide 1197 is "G" in this variant compared to "A" in the Ellison et al sequence (silent mutation)

35 The fusion protein was expressed transiently in Cos1 cells using the pCDN vector system, as described in International Patent Application Publication number WO 96/04388. The mature protein was exported from the cells into the culture medium and was detected by anti-leptin antibody. It was shown to have a size consistent with the predicted structure by Western blotting analysis under both reducing and nonreducing conditions.

40

**Table 7. DNA sequence of ob/IgG1'GT' chimera , 1203bp**

ATGCATTGGGGAACCCTGTGCGGATTCTTGTGGCTTTGGCCCTATCTTTTCTATGTCCAA

45

60

AAG  
1203

5

Table 8. Amino acid sequence of leptin/IgG1 'GT' chimera, 401aa

1 MHWGTLCGFL WLWPYLFYVQ AVPIQKVQDD TKTLIKTIVT  
10 RINDISHTQS  
  
51 VSSKQKVTGL DFIPGLHPIL TLSKMDQTLA VYQQILTSMP SRNVIQISND  
  
101 LENLRDLLHV LAFSKSCHLP WASGLETLDL LGGVLEASGY  
15 STEVVALSRL  
  
151 QGSLQDMLWQ LDLSPGCGTE PKSADKTHTC PPCPAPELLG  
GPSVFLFPPK  
  
20 201 PKDTLMISRT PEVTCVVVDV SHEDPEVKFN WYVDGVEVHN  
AKTKPREEQY  
  
251 NSTYRVVSVL TVLHQDWLNG KEYKCKVSNK ALPAPIEKTI  
SKAKGQPREP  
25 301 QVYTLPPSRE EMTKNQVSLT CLVKGFYPSD IAVEWESNGQ  
PENNYKTPPP  
  
351 VLDSGDGSFFL YSKLTVDKSR WQQGNVFSCS VMHEALHNHY  
30 TQKSLSLSPG  
  
401 K

35

- iii) culturing said transformed host cell under conditions permitting expression of said DNA polymer to produce said chimera; and
- iv) recovering said chimera.

- 5 12. A DNA polymer comprising a nucleotide sequence that encodes a chimera according to any one of claims 1 to 8.
13. A vector which comprises a DNA polymer according to claim 12.
- 10 14. A host cell transformed or transfected with a DNA polymer according to claim 12 or a vector according to claim 13.
- 15 15. A pharmaceutical composition comprising a chimera as claimed in claim 1 and a pharmaceutically acceptable carrier.
16. A chimera according to claim 1, for use as an active therapeutic substance.
17. A chimera according to claim 1, for use in the treatment of obesity or diseases associated with obesity.
- 20 18. A method for the treatment of obesity or diseases associated with in human or non-human mammal, which method comprises administering to the sufferer an effective, non-toxic amount of a chimera as claimed in claim 1.
- 25 19. A chimera as claimed in claim 1, for use in the cosmetic treatment of human or non-human mammals.
20. A method for the cosmetic treatment of a human or non-human mammal, which treatment comprises administering an effective, non-toxic amount of a compound of the
- 30 invention to a human or non-human mammal in need thereof.